

**“JUST THE MATHS”**

**UNIT NUMBER**

**11.1**

**DIFFERENTIATION APPLICATIONS 1**  
**(Tangents and normals)**

by

**A.J.Hobson**

**11.1.1 Tangents**  
**11.1.2 Normals**  
**11.1.3 Exercises**  
**11.1.4 Answers to exercises**

## UNIT 11.1 - APPLICATIONS OF DIFFERENTIATION 1

### TANGENTS AND NORMALS

#### 11.1.1 TANGENTS

In the definition of a derivative (Unit 10.2), it is explained that the derivative of the function  $f(x)$  can be interpreted as the gradient of the tangent to the curve  $y = f(x)$  at the point  $(x, y)$ .

We may now use this information, together with the geometry of the straight line, in order to determine the equation of the tangent to a given curve at a particular point on it.

We illustrate with examples which will then be used also in the subsequent paragraph dealing with normals.

#### EXAMPLES

1. Determine the equation of the tangent at the point  $(-1, 2)$  to the curve whose equation is

$$y = 2x^3 + 5x^2 - 2x - 3.$$

#### Solution

$$\frac{dy}{dx} = 6x^2 + 10x - 2,$$

which takes the value  $-6$  when  $x = -1$ .

Hence the tangent is the straight line passing through the point  $(-1, 2)$  having gradient  $-6$ . Its equation is therefore

$$y - 2 = -6(x + 1).$$

That is,

$$6x + y + 4 = 0.$$

2. Determine the equation of the tangent at the point  $(2, -2)$  to the curve to the curve whose equation is

$$x^2 + y^2 + 3xy + 4 = 0.$$

#### Solution

$$2x + 2y \frac{dy}{dx} + 3 \left[ x \frac{dy}{dx} + y \right] = 0.$$

That is,

$$\frac{dy}{dx} = -\frac{2x + 3y}{3x + 2y},$$

which takes the value  $-2$  at the point  $(2, -2)$ .

Hence, the equation of the tangent is

$$y + 2 = -2(x - 2).$$

That is,

$$2x + y - 2 = 0.$$

3. Determine the equation of the tangent at the point where  $t = 2$  to the curve given parametrically by

$$x = \frac{3t}{1+t} \quad \text{and} \quad y = \frac{t^2}{1+t}.$$

### Solution

We note first that the point at which  $t = 2$  has co-ordinates  $(2, \frac{4}{3})$ .

Furthermore,

$$\frac{dx}{dt} = \frac{3}{(1+t)^2} \quad \text{and} \quad \frac{dy}{dt} = \frac{2t+t^2}{(1+t)^2},$$

by the quotient rule.

Thus,

$$\frac{dy}{dx} = \frac{2t+t^2}{3},$$

which takes the value  $\frac{8}{3}$  when  $t = 2$ .

Hence, the equation of the tangent is

$$y - \frac{4}{3} = \frac{8}{3}(x - 2).$$

That is,

$$3y + 12 = 8x.$$

### 11.1.2 NORMALS

The normal to a curve at a point on it is defined to be a straight line passing through this point and perpendicular to the tangent there.

Using previous work on perpendicular lines (Unit 5.2), if the gradient of the tangent is  $m$ , then the gradient of the normal will be  $-\frac{1}{m}$ .

#### EXAMPLES

In the examples of section 11.1.1, therefore, the normals to each curve at the point given will have equations as follows:

1.

$$y - 2 = \frac{1}{6}(x + 1).$$

That is,

$$6y = x + 13.$$

2.

$$y + 2 = \frac{1}{2}(x - 2).$$

That is,

$$2y = x - 6.$$

3.

$$y - \frac{4}{3} = -\frac{3}{8}(x - 2).$$

That is,

$$24y + 9x = 50.$$


#### Note:

It may occasionally be required to determine the angle,  $\theta$ , between two curves at one of their points of intersection. This is defined to be the angle between the tangents at this point; and, if the gradients of the tangents are  $m_1 = \tan \theta_1$  and  $m_2 = \tan \theta_2$ , then the angle  $\theta \equiv \theta_2 - \theta_1$  and is given by

$$\tan \theta = \frac{\tan \theta_2 - \tan \theta_1}{1 + \tan \theta_2 \tan \theta_1}.$$

That is,

$$\tan \theta = \frac{m_2 - m_1}{1 + m_2 m_1}.$$



### 11.1.3 EXERCISES

1. Determine the equations of the tangent and normal to the following curves at the point given:

(a)

$$8y = x^3 \quad \text{at } (2, 1);$$

(b)

$$y = \frac{e^{2x} \cos x}{(1+x)^3} \quad \text{at } (0, 1).$$

2. The parametric equations of a curve are

$$x = 1 + \sin 2t, \quad y = 1 + \cos t + \cos 2t.$$

Determine the equation of the tangent to the curve at the point for which  $t = \frac{\pi}{2}$ .

3. Determine the equation of the tangent at the point  $(2, 3)$  to the curve whose equation is

$$3x^2 + 2y^2 = 30.$$

4. Determine the equation of the normal at the point  $(-1, 2)$  to the curve whose equation is

$$2xy + 3xy^2 - x^2 + y^3 + 9 = 0.$$

#### 11.1.4 ANSWERS TO EXERCISES

1. (a) The tangent is

$$2y = 3x - 4,$$

and the normal is

$$2x + 3y = 7;$$

(b) The tangent is

$$y = 1 - x,$$

and the normal is

$$y = x + 1.$$

2. The tangent is

$$2y = x - 1.$$

3. The tangent is

$$x + y = 5.$$

4. The normal is

$$x + 9y = 17.$$