

“JUST THE MATHS”

UNIT NUMBER

13.13

**INTEGRATION APPLICATIONS 13
(Second moments of a volume (A))**

by

A.J.Hobson

13.13.1 Introduction

**13.13.2 The second moment of a volume of revolution about
the y -axis**

**13.13.3 The second moment of a volume of revolution about
the x -axis**

13.13.4 Exercises

13.13.5 Answers to exercises

UNIT 13.13 - INTEGRATION APPLICATIONS 13

SECOND MOMENTS OF A VOLUME (A)


13.13.1 INTRODUCTION

Suppose that R denotes a region (with volume V) in space and suppose that δV is the volume of a small element of this region.

Then the “**second moment**” of R about a fixed line, l , is given by

$$\lim_{\delta V \rightarrow 0} \sum_R h^2 \delta V,$$


where h is the perpendicular distance from l of the element with volume δV .



EXAMPLE

Determine the second moment, about its own axis, of a solid right-circular cylinder with height, h , and radius, a .

Solution



In a thin cylindrical shell with internal radius, r , and thickness, δr , all of the elements of volume have the same perpendicular distance, r , from the axis of moments.

Hence the second moment of this shell will be the product of its volume and r^2 ; that is, $r^2(2\pi r h \delta r)$.


The total second moment is therefore given by

$$\lim_{\delta r \rightarrow 0} \sum_{r=0}^{r=a} r^2(2\pi r h \delta r) = \int_0^a 2\pi h r^3 \, dr = \frac{\pi a^4 h}{2}.$$

13.13.2 THE SECOND MOMENT OF A VOLUME OF REVOLUTION ABOUT THE Y-AXIS

Let us consider a region in the first quadrant of the xy -plane, bounded by the x -axis, the lines $x = a$, $x = b$ and the curve whose equation is

$$y = f(x).$$



The volume of revolution of a narrow 'strip', of width, δx , and height, y , (parallel to the y -axis), is a cylindrical 'shell', of internal radius x , height, y , and thickness, δx .

Hence, from the example in the previous section, its second moment about the y -axis is $2\pi x^3 y \delta x$.

Thus, the total second moment about the y -axis is given by

$$\lim_{\delta x \rightarrow 0} \sum_{x=a}^{x=b} 2\pi x^3 y \delta x = \int_a^b 2\pi x^3 y \, dx.$$


Note:

Second moments about the x -axis will be discussed mainly in the next section of this Unit; but we note that, for the volume of revolution, about the x -axis, of a region in the first quadrant bounded by the y -axis, the lines $y = c$, $y = d$ and the curve whose equation is

$$x = g(y),$$

we may reverse the roles of x and y so that the second moment about the x -axis is given by

$$\int_c^d 2\pi y^3 x \, dy.$$



EXAMPLE


Determine the second moment, about a diameter, of a solid sphere with radius a .

Solution

We may consider, first, the volume of revolution about the y -axis of the region, bounded in the first quadrant, by the x -axis, the y -axis and the circle whose equation is

$$x^2 + y^2 = a^2,$$

then double the result obtained.



The total second moment is given by

$$2 \int_0^a 2\pi x^3 \sqrt{a^2 - x^2} dx = 4\pi \int_0^{\frac{\pi}{2}} a^3 \sin^3 \theta \cdot a \cos \theta \cdot a \cos \theta d\theta,$$

if we substitute $x = a \sin \theta$.

This simplifies to

$$4\pi a^5 \int_0^{\frac{\pi}{2}} \sin^3 \theta \cos^2 \theta d\theta = 4\pi \int_0^{\frac{\pi}{2}} (\cos^2 \theta - \cos^4 \theta) \sin \theta d\theta,$$

if we make use of the trigonometric identity

$$\sin^2 \theta \equiv 1 - \cos^2 \theta.$$


The total second moment is now given by

$$4\pi a^5 \left[-\frac{\cos^3 \theta}{3} + \frac{\cos^5 \theta}{5} \right]_0^{\frac{\pi}{2}} = 4\pi a^5 \left(\frac{1}{3} - \frac{1}{5} \right) = \frac{8\pi a^5}{15}.$$

13.13.3 THE SECOND MOMENT OF A VOLUME OF REVOLUTION ABOUT THE X-AXIS

In the introduction to this Unit, a formula was established for the second moment of a solid right-circular cylinder about its own axis. This result may now be used to determine the second moment about the x -axis for the volume of revolution about this axis of a region enclosed in the first quadrant by the x -axis, the lines $x = a$, $x = b$ and the curve whose equation is

$$y = f(x).$$



The volume of revolution about the x -axis of a narrow strip, of width δx and height y , is a cylindrical 'disc' whose second moment about the x -axis is $\frac{\pi y^4 \delta x}{2}$. Hence the second moment of the whole region about the x -axis is given by


$$\lim_{\delta x \rightarrow 0} \sum_{x=a}^{x=b} \frac{\pi y^4}{2} \delta x = \int_a^b \frac{\pi y^4}{2} dx.$$

EXAMPLE

Determine the second moment about the x -axis for the volume of revolution about this axis of the region bounded in the first quadrant by the x -axis, the y -axis, the line $x = 1$ and the line whose equation is

$$y = x + 1.$$

Solution



$$\text{Second moment} = \int_0^1 \frac{\pi(x+1)^4}{2} dx = \left[\pi \frac{(x+1)^4}{10} \right]_0^1 = \frac{31\pi}{10}.$$

Note:

The second moment of a volume about a certain axis is closely related to its “**moment of inertia**” about that axis. In fact, for a solid, with uniform density, ρ , the Moment of Inertia is ρ times the second moment of volume, since multiplication by ρ of elements of volume converts them into elements of mass.

13.13.4 EXERCISES

1. Determine the second moment about a diameter of a circular disc with small thickness, t , and radius, a .
2. Determine the second moment, about the axis specified, for the volume of revolution of each of the following regions of the xy -plane about this axis:
 - (a) Bounded in the first quadrant by the x -axis, the y -axis and the curve whose equation is

$$y = 1 - 2x^2.$$

Axis: The y -axis.

- (b) Bounded in the first quadrant by the x -axis and the curve whose equation is

$$y^2 = \sin x.$$

Axis: The x -axis.

- (c) Bounded in the first quadrant by the x -axis, the y -axis, the line $x = 1$ and the curve whose equation is

$$y = e^{-2x}.$$

Axis: The x -axis

- (d) Bounded in the first quadrant by the x -axis, the y -axis, the line $x = 1$ and the curve whose equation is

$$y = e^{-2x}.$$

Axis: The y -axis.

13.13.5 ANSWERS TO EXERCISES

1.

$$\frac{\pi a^4 t}{4}.$$

2. (a)

$$\frac{\pi}{24}.$$

(b)

$$\frac{\pi^2}{4}.$$

(c)

0.196, approximately.

(d)

0.337, approximately.