

“JUST THE MATHS”

SLIDES NUMBER

10.6

DIFFERENTIATION 6
(Inverse trigonometric functions)

by

A.J.Hobson

10.6.1 Summary of results
10.6.2 The derivative of an inverse sine
10.6.3 The derivative of an inverse cosine
10.6.4 The derivative of an inverse tangent

UNIT 10.6 - DIFFERENTIATION 6

DERIVATIVES OF INVERSE TRIGONOMETRIC FUNCTIONS

10.6.1 SUMMARY OF RESULTS

1.

$$\frac{d}{dx}[\sin^{-1}x] = \frac{1}{\sqrt{1-x^2}},$$

where $-\frac{\pi}{2} \leq \sin^{-1}x \leq \frac{\pi}{2}$.

2.

$$\frac{d}{dx}[\cos^{-1}x] = -\frac{1}{\sqrt{1-x^2}},$$

where $0 \leq \cos^{-1}x \leq \pi$.

3.

$$\frac{d}{dx}[\tan^{-1}x] = \frac{1}{1+x^2},$$

where $-\frac{\pi}{2} \leq \tan^{-1}x \leq \frac{\pi}{2}$.

10.6.2 THE DERIVATIVE OF AN INVERSE SINE

We shall consider the formula

$$y = \text{Sin}^{-1}x$$

and determine an expression for $\frac{dy}{dx}$.

Note:

There is a special significance in using the upper-case S in the formula (see later).

The formula is equivalent to

$$x = \sin y,$$

so that

$$\frac{dx}{dy} = \cos y \equiv \pm\sqrt{1 - \sin^2 y} \equiv \pm\sqrt{1 - x^2}.$$


Thus,

$$\frac{dy}{dx} = \pm\frac{1}{\sqrt{1 - x^2}}.$$

Consider now the graph of the formula


$$y = \text{Sin}^{-1}x.$$

This may be obtained from the graph of $y = \sin x$ by reversing the roles of x and y and rearranging the new axes into the usual positions.



Observations

- (i) x must lie in the interval $-1 \leq x \leq 1$.
- (ii) For each x in $-1 \leq x \leq 1$, y has infinitely many values - spaced at regular intervals of $\frac{\pi}{2}$.
- (iii) For each x in the interval $-1 \leq x \leq 1$, there are only two possible values of $\frac{dy}{dx}$, one positive and the other negative.



(iv) On the part of the graph from $y = -\frac{\pi}{2}$ to $y = \frac{\pi}{2}$, there will be only one value of y and one (positive) value of $\frac{dy}{dx}$ for each x in $-1 \leq x \leq 1$.

The restricted part of the graph defines the “**principal value**” of the inverse sine function and is denoted by $\sin^{-1}x$ using a lower-case s.

Hence,

$$\frac{d}{dx}[\sin^{-1}x] = \frac{1}{\sqrt{1-x^2}}.$$

10.6.3 THE DERIVATIVE OF AN INVERSE COSINE

We shall consider the formula

$$y = \text{Cos}^{-1}x$$

and determine an expression for $\frac{dy}{dx}$.

Note:

There is a special significance in using the upper-case C in the formula (see later).

The formula is equivalent to

$$x = \cos y,$$

so that

$$\frac{dx}{dy} = -\sin y \equiv \pm\sqrt{1 - \cos^2 y} \equiv \pm\sqrt{1 - x^2}.$$


Thus,

$$\frac{dy}{dx} = \pm\frac{1}{\sqrt{1 - x^2}}.$$

Consider now the graph of the formula


$$y = \text{Cos}^{-1}x,$$

which may be obtained from the graph of $y = \cos x$ by reversing the roles of x and y and rearranging the new axes into the usual positions.



Observations

- (i) x must lie in the interval $-1 \leq x \leq 1$.
- (ii) For each x in $-1 \leq x \leq 1$, y has infinitely many values - spaced at regular intervals of $\frac{\pi}{2}$.
- (iii) For each x in the interval $-1 \leq x \leq 1$, there are only two possible values of $\frac{dy}{dx}$, one positive and the other negative.



(iv) On the part of the graph from $y = 0$ to $y = \pi$, we may distinguish the results from those of the inverse sine function.

There will be only one value of y with one (negative) value of $\frac{dy}{dx}$ for each x in $-1 \leq x \leq 1$.

The restricted part of the graph defines the “**principal value**” of the inverse cosine function and is denoted by $\cos^{-1}x$ using a lower-case c.

Hence,

$$\frac{d}{dx}[\cos^{-1}x] = -\frac{1}{\sqrt{1-x^2}}.$$

10.6.4 THE DERIVATIVE OF AN INVERSE TANGENT

We shall consider the formula

$$y = \text{Tan}^{-1}x$$

and determine an expression for $\frac{dy}{dx}$.

Note:

There is a special significance in using the upper-case T in the formula (see later).

The formula is equivalent to

$$x = \tan y,$$

so that

$$\frac{dx}{dy} = \sec^2 y \equiv 1 + \tan^2 y \equiv 1 + x^2.$$


Thus,

$$\frac{dy}{dx} = \frac{1}{1 + x^2}.$$

Consider now the graph of the formula


$$y = \text{Tan}^{-1}x,$$

which may be obtained from the graph of $y = \tan x$ by reversing the roles of x and y and rearranging the new axes into the usual positions.



Observations

- (i) x may lie anywhere in the interval $-\infty < x < \infty$.
- (ii) For each x , y has infinitely many values - spaced at regular intervals of π .
- (iii) For each x , there is only possible value of $\frac{dy}{dx}$, which is positive.



(iv) On the part of the graph from $y = -\frac{\pi}{2}$ to $y = \frac{\pi}{2}$, there will be only one value of y for each value of x .

The restricted part of the graph defines the “**principal value**” of the inverse tangent function and is denoted by $\tan^{-1}x$ using a lower-case t.

Hence,

$$\frac{d}{dx}[\tan^{-1}x] = \frac{1}{1+x^2}.$$

ILLUSTRATIONS

1.

$$\frac{d}{dx}[\sin^{-1}2x] = \frac{2}{\sqrt{1-4x^2}}.$$

2.

$$\frac{d}{dx}[\cos^{-1}(x+3)] = -\frac{1}{\sqrt{1-(x+3)^2}}.$$

3.

$$\frac{d}{dx}[\tan^{-1}(\sin x)] = \frac{\cos x}{1 + \sin^2 x}.$$

4.

$$\frac{d}{dx}[\sin^{-1}(x^5)] = \frac{5x^4}{\sqrt{1 - x^{10}}}.$$