

**“JUST THE MATHS”**

**SLIDES NUMBER**

**2.2**

**SERIES 2  
(Binomial series)**

by

**A.J.Hobson**

2.2.1 Pascal's Triangle  
2.2.2 Binomial Formulae

## UNIT 2.2 - SERIES 2 - BINOMIAL SERIES

### INTRODUCTION

In this section, we expand (multiplying out) an expression of the form

$$(A + B)^n.$$

$A$  and  $B$  can be either mathematical expressions or numerical values.

$n$  is a given number which need not be a positive integer.

### 2.2.1 PASCAL'S TRIANGLE

#### ILLUSTRATIONS

1.  $(A + B)^1 \equiv$

$$A + B.$$

2.  $(A + B)^2 \equiv$

$$A^2 + 2AB + B^2.$$

3.  $(A + B)^3 \equiv$

$$A^3 + 3A^2B + 3AB^2 + B^3.$$

4.  $(A + B)^4 \equiv$

$$A^4 + 4A^3B + 6A^2B^2 + 4AB^3 + B^4.$$

#### OBSERVATIONS

(i) The expansions begin with the maximum possible



$$6. (A - B)^6 \equiv$$

$$A^6 - 6A^5B + 15A^4B^2 - 20A^3B^3 + 15A^2B^4 - 6AB^5 + B^6.$$

### 2.2.2 BINOMIAL FORMULAE

A more general method which can be applied to any value of  $n$  is the binomial formula.

#### DEFINITION

If  $n$  is a positive integer, the product

$$1.2.3.4.5.....n$$

is denoted by the symbol  $n!$  and is called “ $n$  factorial”.

#### Note:

This definition could not be applied to the case when  $n = 0$ .

$0!$  is defined separately by the statement

$$0! = 1.$$

There is no meaning to  $n!$  when  $n$  is a negative integer.

**(a) Binomial formula for  $(A + B)^n$  when  $n$  is a positive integer.**

It can be shown that

$$(A + B)^n \equiv A^n + nA^{n-1}B + \frac{n(n-1)}{2!}A^{n-2}B^2 + \frac{n(n-1)(n-2)}{3!}A^{n-3}B^3 + \dots + B^n.$$

**Notes:**

(i) This is the same result as given by Pascal's Triangle.

(ii) The last term is

$$\frac{n(n-1)(n-2)(n-3)\dots\dots 3.2.1}{n!}A^{n-n}B^n = A^0B^n = B^n.$$

(iii) The coefficient of  $A^{n-r}B^r$  in the expansion is

$$\frac{n(n-1)(n-2)(n-3)\dots\dots(n-r+1)}{r!} = \frac{n!}{(n-r)!r!}$$

and this is sometimes denoted by the symbol  $\binom{n}{r}$ .

(iv) A commonly used version is

$$(1 + x)^n \equiv 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3 + \dots + x^n.$$

## EXAMPLES

1. Expand fully the expression  $(1 + 2x)^3$ .

**Solution**

$$(A + B)^3 \equiv A^3 + 3A^2B + 3AB^2 + B^3.$$

Replace  $A$  by 1 and  $B$  by  $2x$ .

$$\begin{aligned}(1 + 2x)^3 &\equiv 1 + 3(2x) + 3(2x)^2 + (2x)^3 \\ &\equiv 1 + 6x + 12x^2 + 8x^3.\end{aligned}$$

2. Expand fully the expression  $(2 - x)^5$ .

**Solution**

$$(A + B)^5 \equiv A^5 + 5A^4B + 10A^3B^2 + 10A^2B^3 + 5AB^4 + B^5.$$

Replace  $A$  by 2 and  $B$  by  $-x$ .

$$\begin{aligned}(2 - x)^5 &\equiv 2^5 + 5(2)^4(-x) + 10(2)^3(-x)^2 + \\ &\quad 10(2)^2(-x)^3 + 5(2)(-x)^4 + (-x)^5.\end{aligned}$$

That is,

$$(2 - x)^5 \equiv 32 - 80x + 80x^2 - 40x^3 + 10x^4 - x^5.$$

**(b) Binomial formula for  $(A + B)^n$  when  $n$  is negative or a fraction.**

This time, the series will be an **infinite** series.

## RESULT

If  $n$  is negative or a fraction and  $x$  lies strictly between  $x = -1$  and  $x = 1$ , it can be shown that

$$(1 + x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3 + \dots$$

## EXAMPLES

1. Expand  $(1 + x)^{\frac{1}{2}}$  as far as the term in  $x^3$ .

**Solution**

$$(1 + x)^{\frac{1}{2}} = 1 + \frac{1}{2}x + \frac{\frac{1}{2}(\frac{1}{2} - 1)}{2!}x^2 + \frac{\frac{1}{2}(\frac{1}{2} - 1)(\frac{1}{2} - 2)}{3!}x^3 + \dots$$

$$= 1 + \frac{x}{2} - \frac{x^2}{8} + \frac{x^3}{16} - \dots$$

provided  $-1 < x < 1$ .

2. Expand  $(2 - x)^{-3}$  as far as the term in  $x^3$  stating the values of  $x$  for which the series is valid.

**Solution**

First convert the expression  $(2 - x)^{-3}$  to one in which the leading term in the bracket is 1.

$$\begin{aligned}(2 - x)^{-3} &\equiv \left[2 \left(1 - \frac{x}{2}\right)\right]^{-3} \\ &\equiv \frac{1}{8} \left(1 + \left[-\frac{x}{2}\right]\right)^{-3}.\end{aligned}$$

The required binomial expansion is

$$\begin{aligned}\frac{1}{8} &\left[1 + (-3) \left(-\frac{x}{2}\right) + \frac{(-3)(-3-1)}{2!} \left(-\frac{x}{2}\right)^2 + \right. \\ &\left. \frac{(-3)(-3-1)(-3-2)}{3!} \left(-\frac{x}{2}\right)^3 + \dots\right].\end{aligned}$$

That is,

$$\frac{1}{8} \left[1 + \frac{3x}{2} + \frac{3x^2}{2} + \frac{5x^3}{4} + \dots\right].$$

The expansion is valid provided  $-x/2$  lies strictly between  $-1$  and  $1$ .

Hence,  $-2 < x < 2$ .

### (c) Approximate Values

The Binomial Series may be used to calculate simple approximations, as illustrated by the following example:

#### EXAMPLE

Evaluate  $\sqrt{1.02}$  correct to five places of decimals.

#### Solution

Using  $1.02 = 1 + 0.02$ , we may say that

$$\sqrt{1.02} = (1 + 0.02)^{\frac{1}{2}}.$$

That is,

$$\sqrt{1.02} = 1 + \frac{1}{2}(0.02) + \frac{\frac{1}{2}(-\frac{1}{2})}{1.2}(0.02)^2 + \frac{\frac{1}{2}(-\frac{1}{2})(-\frac{3}{2})}{1.2.3}(0.02)^3 + \dots$$

$$= 1 + 0.01 - \frac{1}{8}(0.0004) + \frac{1}{16}(0.000008) - \dots$$

$$= 1 + 0.01 - 0.00005 + 0.0000005 - \dots$$

$$\simeq 1.010001 - 0.000050 = 1.009951$$

Hence,  $\sqrt{1.02} \simeq 1.00995$